




Nance Appliance



The Nance appliance is a very effective retainer that prevents the back teeth from moving forward. It is important that an extra effort be made to clean the appliance on the roof of the mouth. This will prevent the gums around the Nance appliance from getting inflamed and swollen. Speech will return to normal after the first few days. In order to prevent irritation during sleep, you may place wax on the cheek-side of the teeth that have metal bands.

You should not drink any carbonated beverage as the carbonation can dissolve the cement used to cement bands to the teeth. We also ask that you NOT eat anything which fits into the category of "hard, sticky or chewy." Below we have listed some foods which we would like for you to avoid. **This list does not include all foods to avoid.** It is only a guideline to help you become more selective about the foods to be eaten to prevent damage to the appliance. In addition to these foods, **pens, pencils, ice or anything hard** which you put into your mouth can also damage the appliance or your teeth.

Hard Foods:	Popcorn, Nuts, Peanut brittle, Hard taco shells, Doritos, Nachos, Pizza crust, Breath Mints, Ice, etc.
Sticky/Chevy Foods:	Caramel, Chewing Gum, Bubble Gum, Taffy, Jolly Ranchers, Skittles, Gummy Bears, Licorice, etc.


There are still many things that you CAN eat with your orthodontic appliance. For example, fruit, raisins, Hershey Kisses, Nestle Crunch bars, Three Musketeers, M&M's (not peanut or almond), Reese's Peanut Butter Cups or Pieces, Peppermint Patties, Cotton Candy, Ice Cream, Frozen Yogurt, Brownies (without nuts), etc.

It is a good idea to brush your teeth soon after eating a meal or snack to remove the food from around your teeth, gums and appliance. If you are unable to brush, then rinse well with water.

Please feel free to call us with any questions you may have!



Aparato Nance



El aparato Nance es un retenedor muy efectivo que evita que los dientes posteriores se muevan hacia adelante.

Es importante que se haga un esfuerzo adicional para limpiar el aparato en el techo de la boca. Esto evitará que las encías alrededor del aparato Nance se inflamen o hinchen.

El habla volverá a la normalidad después de los primeros días. Para evitar la irritación mientras duerme,

puede colocar cera en el lado de la mejilla de los dientes que tienen bandas de metal.

No debe beber ninguna bebida carbonatada ya que la carbonatación puede disolver el cemento utilizado para en las bandas de los dientes. También le pedimos que NO coma nada que se ajuste a la categoría de "duro, pegajoso o masticable". A continuación están algunos alimentos que nos gustaría evitar. Esta lista no incluye todos los alimentos. Es solo una guía para ayudarlo a ser más selectivo acerca de los alimentos que se deben comer para evitar daños al retenedor. Además de estos alimentos, lapiceros, lápices, hielo o cualquier cosa dura que se ponga en la boca también puede dañar el retenedor o los dientes.

Alimentos duros:	Palomitas de maíz, nueces, tacos duros, doritos, nachos, pan de la pizza, mentas para aliento, hielo, etc.
Alimentos pegajosos / masticables:	Caramelo, chicle, Jolly Ranchers, skittles, ositos de goma, etc.

Todavía hay muchas cosas que PUEDE comer con su aparato de ortodoncia. Por ejemplo, fruta, pasas, chocolate Hershey, barritas Nestle Crunch, tres mosqueteros, M&M (no cacahuate o almendra), empanadas de menta, algodón de azúcar, helado, yogurt, brownies (sin nueces), etc.

Es una buena idea cepillarse los dientes poco después de comer para quitar la comida de los dientes, encías y del retenedor. Si no puede cepillarse, enjuague bien con agua.

¡No dude en llamarnos si tiene alguna pregunta!